



Real history of helical vortices: theory and applications. Annex: comments on “motion of a helical vortex by o. Velasco fuentes”

Okulov, Valery; Sørensen, Jens Nørkær; Fukumoto, Yasuhide; Wood, David H.

Publication date:
2018

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Okulov, V., Sørensen, J. N., Fukumoto, Y., & Wood, D. H. (2018). *Real history of helical vortices: theory and applications. Annex: comments on “motion of a helical vortex by o. Velasco fuentes”*. 71-72. Abstract from 4th International Retreat on Vortical Flow and Aerodynamics (IRVA4) , Novosibirsk, Russian Federation.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

BOOK OF ABSTRACTS

4th International Retreat on Vortical Flow and Aerodynamics

Novosibirsk, Russia, October 1 – 3, 2018

Honorary Chairs: **Professor James Wu**
Shanghai Jiaotong University, Shanghai, China

Professor Sergey Alekseenko
*Kutateladze Institute of Thermophysics, SB RAS,
Novosibirsk, Russia*

Co-Chairs: **Professor Peng Bai**
*China Academy of Aerospace Aerodynamics,
Beijing, China*

Professor Valery Okulov
Technical University of Denmark, Denmark

Professor Jens Sørensen
Technical University of Denmark, Denmark

Colloquium Organizers:

- China Academy of Aerospace Aerodynamics
- Technical University of Denmark
- Russian Academy of Sciences
- Kutateladze Institute of Thermophysics, SB RAS
- Khristianovich Institute of Theoretical and Applied Mechanics, SB RAS

Sponsor:

- Russian Foundation for Basic Research

Website: <http://www.itp.nsc.ru/konferentsiya-irva4/konferentsiya>

REAL HISTORY OF HELICAL VORTICES: THEORY AND APPLICATIONS.

Annex: Comments on “Motion of a helical vortex by O. Velasco Fuentes”

Valery L. Okulov^{1,2}, Jens N. Sørensen¹, Yasuhide Fukumoto³, David H. Wood⁴

¹ Department of Wind Energy, Technical University of Denmark, Lyngby, Denmark

² Kutateladze Institute of Thermophysics, SB RAS, Novosibirsk, Russia

³ Institute of Mathematics for Industry, Kyushu University, Fukuoka, Japan

⁴ Schulich School of Engineering, University of Calgary Canada

E-mail: vokulov@mail.ru

Keywords: helical vortex, vortex precession, self-induced motion

Research on helical vortices has a long history, which dates back to the famous work of Lord Kelvin in 1880 on the helical perturbations of the columnar vortex. Helical vortices are of fundamental importance to fluid mechanics because they describe one of the main states of swirling flows. An accurate estimation of velocities in the rotor plane requires knowing the velocity field induced by the helical vortices. Therefore, starting with the inspirational work of Joukowski, the theory of helical vortices has been actively studied as a prerequisite to understand and analyze rotor aerodynamics [1]. At present the fundamentals are based on various analytical components [1, 2], which holds true for all values of the helix pitch, such as (i) the 2D Biot-Savart law for helical filaments represented by Kapteyn series or in a form with singularity separation; (ii) solutions of helical vortex tubes with finite core, governed by series expansion of helical multipoles; (iii) relations between the induction of vortex filaments and the self-induced velocity of helical vortex tubes resulting in a closed analytical solution of the helix motion; (iv) analytical representation of Goldstein’s solution for the circulation of a helical vortex sheet in equilibrium; (v) Kelvin’s N-gon stability problem of point vortices generalized to multiple helical vortices.

Annex: Comments on “Motion of a helical vortex by O. Velasco Fuentes”

In the recent paper on the motion of helical vortices [VF2018] in J. Fluid Mech. 836, R1, doi:10.1017/jfm.2017.845 the model with both tangential and binormal velocities including in the helical motion was used in contrast to the classical approach when only the bi-normal component included [3]. A large difference between both solutions was found. We demonstrate here that conflicts of VF2018 were based on a bias in the representation of the velocities in different coordinate systems and that the motion of a helical vortex is uniquely dictated by the bi-normal component of the velocity as it was stated in the classical approach and only it correlates with the Betz-Joukowski limit well [1].




Figure 1: Correlations of azimuthal velocities u_θ inducing by infinite thin vortex line (dotted lines) and vortex with the finite cores of uniform vorticity ω_t distributions (solid lines):
(a) rectilinear vortices and (b) helical vortices [4].

VF2018 also erroneously expand a property of the point vortex dynamics to the helical vortex (Fig. 1). He used the vortex-line solutions (dotted lines) instead of one for the finite core (solid lines) for the helical vortex with a finite core in which even the uniform vorticity concludes to non-uniform velocity profiles analytically proven by Eq. (3.6) of [5].

A comparison of the Ω of [3] with an axial motion of fluid particles along the helix axis

The main error of the VF2018 is his comparison of the angular velocities (red VF2018 and green [3] curves on fig 2) because both ones were derived in different coordinate systems. The vortex system in [3] p. 328, 335 moves with the fixed velocity $\Gamma N/(2\pi R\tau)$. For a partial case (Fig. 2) when the fluid particles move with the u_a only in the axial direction we will show that the classical approach of [3] can explain both axial u_a and azimuthal $\Delta\theta/\Delta t$ displacement of the helical vortex when the VF2018 gives only one of that, u_a (or $\Omega^*(\tau_0) = 0$).




Figure 2: Sketch of the helix displacement with the axial translation of the fluid particles u_a and the plots of the angular and axial displacements of the helix (green lines - [3]) and fluid particles (red lines - VF2018).

The root of red curve of VF2018 gives the $\tau_0 \approx 0.275$ and the correspondence value of the absolute velocity in this point $u_a(\tau_0) \approx U^*(\tau_0) = 5.35$. The angular velocity of [3] (green curve) has non zero value $\Omega_{Ok}(\tau_0) \approx 0.554$ at $\tau_0 = 0.275$. The coordinate system for the single vortex in [3] p. 328, 335, moves with the fixed velocity $\Gamma/(2\pi R\tau_0) = 4\pi/(2\pi\tau_0) = 2/\tau_0$. If we put the coordinate translation it in the formula $U_{Ok} = (\Gamma/(2\pi R) - R\Omega_{Ok}(\tau_0))/\tau_0$ of [3] for the axial velocity the value $U_{Ok} \approx (2 - 0.554)/0.275 = 5.28$. The correction to the non-uniform flow (fig 1b) with add-on $\Delta\Omega = 0.25\tau/(1+\tau^2)^{3/2} = 0.25 \cdot 0.75/(1+0.75^2)^{3/2} = 0.062$. Summing both lines: $5.28 + 0.062 = 5.342$ which coincides with the axial movement of the fluid particles $U^*(\tau_0) = 5.35$. So the classical approach of [3] permits to estimate both real displacements of the helix in axial, $U_{Ok} \equiv u_a$, and azimuthal, $\Delta\theta/\Delta t = \Omega_{Ok}(\tau_0)$, directions when the by fluid-particle motion of VF2018 approximates only the axial displacement $U^* \approx u_a$ without rotation $\Omega^* = 0$.

Acknowledgements

The work was supported in part by Grant JSPS no. S17119.

References

- [1] Okulov, V.L., Sørensen J.N., & Wood D.H., The rotor theories by Professor Joukowsky, Vortex Theories, *Prog. Aerosp. Sci.*, 2015, 73, pp. 19-46. DOI: 10.1016/j.paerosci.2014.10.002
- [2] Okulov, V.L., & Sørensen, J.N., Applications of 2D helical vortex dynamics, *Theor. Comput. Fluid Dyn.*, 2010, 24, 395-401.
- [3] V.L. Okulov. On the stability of multiple helical vortices. *J. Fluid Mech.* 2004, 521, 319–342.
- [4] Y. Fukumoto & V.L. Okulov. The Velocity Field Induced by a helical Vortex Tube. *Phys. Fluids*, 2005, 17(10), 107101.
- [5] J. Boersma & D. H. Wood. On the self-induced motion of a helical vortex. *J. Fluid Mech.* 1999, 384, 263–280.